



# XENA<sup>®</sup>

## Spreader-Sticker

A Biodegradable, Natural Spreader & Sticker  
For All Crops.

### Keep Out of Reach of Children

### WARNING

May be harmful if swallowed.

**Precautionary Statements:** Do not ingest. Do not get in eyes, skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking or chewing gum.

See back panel for additional precautionary information.

### Shake well before using

Manufactured by:

SAN Group Biotech USA, Inc.  
1260 Avenida Chelsea, Vista, CA 92081 USA

WASDA Reg. No. 51517-22004 • CA Reg. No. 51517-50005-AA



### Principal Functioning Agents:

Polyoxyethylene Sorbitan Monolaurate,  
Soapbark (Quillaja Saponin), Ethoxylated  
Alcohol, Sodium Carboxymethyl

Cellulose.....18.0%

### Constituents Ineffective As Spray

Adjuvants:.....82.0%

TOTAL:..... 100.0%

**Net Content:**  
**2.5 Gallons (9.46 L)**

# XENA<sup>®</sup>

## Spreader-Sticker

### PRODUCT INFORMATION

**XENA<sup>®</sup> Spreader-Sticker** is an environmentally-friendly, biodegradable, natural adherent designed to increase the spreading and adhesion of fungicides, insecticides, miticides, herbicides and desiccants to the plant. Due to its dispersing capacity, **XENA<sup>®</sup> Spreader-Sticker** allows for a homogenous distribution of the applied chemicals. **XENA<sup>®</sup> Spreader-Sticker** improves the effectiveness of fungicides, insecticides, miticides, herbicides and desiccants by reducing the wash off by rain or irrigation

**XENA<sup>®</sup> Spreader-Sticker** is not for aquatic use.

### KEEP OUT OF REACH OF CHILDREN

### WARNING

May be harmful if swallowed.

**PRECAUTIONARY STATEMENT:** Do not ingest. Do not get on skin or clothes. Wash thoroughly with soap and water after handling and before eating, drinking or chewing gum.

### FIRST AID

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything to an unconscious person.

### STORAGE AND DISPOSAL

Store out of direct sunlight and at temperatures above freezing (0°C). Triple rinse and offer empty container for recycling or reconditioning or dispose of in a sanitary landfill.

**NOTICE OF WARRANTY – READY CAREFULLY.** SAN Group Biotech USA, Inc., warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. The directions for use of this product are believed to be reliable and should be followed carefully. It is impossible to eliminate all risks associated with use. Buyer assumes all risks associated with use of this product contrary to label instructions or resulting from extraordinary conditions. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. In no case shall SAN Group Biotech USA, Inc., be liable for special, indirect or consequential damages resulting from the use or handling of this product and no claim of any kind shall be greater than the purchase price of the product in respect of which such damages are claimed. SAN GROUP BIOTECH USA, INC., MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

## DIRECTIONS FOR USE

**XENA<sup>®</sup> Spreader-Sticker** can be applied to all crops including fruit trees, vines, tree nuts, fruits and vegetables, field and row crops, ornamental trees, shrubs and flowers, containerized plants, and turf. Always perform a phytotoxicity test on a small plot before large-scale application. To lower the risk of phytotoxicity, applications should be made early morning or late evening.

**Compatibility:** When mixing **XENA<sup>®</sup> Spreader-Sticker** with any other products read and follow the precautions, restrictions and recommendations on all product labels. A standard jar compatibility test should be conducted before using any new mixes. **DO NOT apply this product when temperatures are above 85°F (29°C) with Copper and/or Sulfur in the tank mix.**

### RECOMMENDED USE RATES:

Always add **XENA<sup>®</sup> Spreader-Sticker** last, after all other products have been added, when spray tank is ¾ full and with agitation.

### Ground or Aerial Applications:

For ground application use up to 4 fluid ounces per 100 gallons of water. For aerial application and for SUPPRESS Herbicide use up to 8 fluid ounces per 100 gallons of water. Follow side chart for rates.

**Nursery & Container:** 1 teaspoon (5 milliliters) per 5 gallons of water.

**Home & Garden:** 1 teaspoon (5 milliliters) per 5 gallons of water.

For other applications refer to the Suggested Use Rates for **XENA<sup>®</sup> Spreader-Sticker** per Gallons of Spray Mix table on the right.

Suggested Use Rates  
per Gallons of Spray Mix

Gallon of Spray Mix	Oz of Spreader-Sticker
50	2
100	4
150	6
200	8
250	10
300	12
400+	16

# SAN<sup>®</sup>

## AGROW

**Manufactured by:**  
**SAN Group Biotech USA, Inc.**  
1260 Avenida Chelsea, Vista,  
CA 92081 USA • Tel. (800) 876-2767  
[www.san-agrow.com](http://www.san-agrow.com)



### NET CONTENTS/NET WEIGHT:

■ 2.5 US Gallons (9.45 L) / 21.5 Lbs (9.76 Kg)

**WASDA Reg. No. 51517-22004 - CA Reg. No. 51517-50005-AA**

©XENA and SUPPRESS are registered trademarks of SAN Group Biotech USA, Inc.